

Sphaleron rate in the symmetric electroweak phase

Guy D. Moore

Department of Physics, University of Washington, Seattle WA 98195-1560 USA

(January 2000)

Abstract

Recently Bödeker has presented an effective infrared theory for the dynamics of Yang-Mills theory, suitable for studying the rate of baryon number violation in the early universe. We extend his theory to include Higgs fields, and study how much the Higgs affects the baryon number violation rate in the symmetric phase, at the phase coexistence temperature of a first order electroweak phase transition. The rate is about 20% smaller than in pure Yang-Mills theory. We also analyze the sphaleron rate in the analytic crossover regime.

I. INTRODUCTION

Baryon number is violated in the standard model [1], and while the rate of the violation is negligibly small in vacuum, it can be significant at higher temperatures [2,3].

Recently there has been substantial interest in how fast baryon number is violated in the standard model at high temperatures. This is both because the rate is important for baryogenesis, and because it is an interesting problem in the more general framework of understanding dynamics of hot gauge theories.

The last 3 years have seen major progress in this problem. First, analytic work has clarified what physics is relevant. While it has long been known that infrared or “soft” gauge fields with momenta $p \sim g^2 T$ are responsible for baryon number violation [3], the recent work of Arnold, Son, and Yaffe has shown that “hard” modes with momenta $p \sim T$ play an essential role in modifying the dynamics of the $p \sim g^2 T$ degrees of freedom [4–6]. Bödeker has gone further, showing the role of scatterings between such hard modes by exchange of $g^2 T \lesssim p \lesssim gT$ modes [7–9]. There has also been progress in numerically studying the effective infrared (IR) theories to determine the size of baryon number violation [10,11].

The rate of baryon number violation is characterized by the sphaleron rate¹

$$\Gamma \equiv \lim_{V, t \rightarrow \infty} \frac{\langle (N_{\text{CS}}(t) - N_{\text{CS}}(0))^2 \rangle}{Vt}, \quad (1.1)$$

$$N_{\text{CS}}(t) - N_{\text{CS}}(0) = \frac{1}{16\pi^2} \int_0^t \int d^3x \epsilon_{ijk} E_i^a F_{jk}^a. \quad (1.2)$$

¹Our normalization is $D_i = \partial_i + iT^a A_i^a$, and $F_{ij} = [D_i, D_j]$. α means α_w , and g means g_w .

Bödeker has shown that in pure Yang-Mills theory, Γ has the parametric form [7]

$$\Gamma = \kappa' \left(\log \frac{m_D}{g^2 T} + C + O(1/\log) \right) \left(\frac{g^2 T^2}{m_D^2} \right) \alpha^5 T^4. \quad (1.3)$$

The constant² C can and recently has been determined analytically [12]. The leading coefficient κ' can be obtained numerically by studying a local, UV finite theory,

$$\sigma D_\tau A_i^a(x, \tau) = -g^2 \frac{\partial H_A(A(\tau))}{\partial A_i^a(x, \tau)} + \xi_i^a(x, \tau), \quad (1.4)$$

$$H_A = \int d^3x \frac{1}{4g^2} F_{ij}^a F_{ij}^a(x), \quad -g^2 \frac{\partial H_A}{\partial A_i^a} = (D_j F_{ji})^a = (D \times B)_i^a, \quad (1.5)$$

$$\langle \xi_i^a(x_1, \tau_1) \xi_j^b(x_2, \tau_2) \rangle = 2g^2 T \delta(x_1 - x_2) \delta(\tau_1 - \tau_2) \delta_{ij} \delta^{ab}, \quad (1.6)$$

with D_τ the covariant derivative in Langevin time τ , F_{ij}^a the (nonabelian) magnetic field, and ξ Gaussian white noise normalized as shown.³ κ' is determined in this theory as

$$\kappa' = \frac{3\sigma}{2\pi} \lim_{V, \tau \rightarrow \infty} \frac{\langle ((N_{CS}(\tau) - N_{CS}(0))^2) \rangle}{V\tau}. \quad (1.7)$$

This effective theory was studied numerically in [13], with the result (see below) that $\kappa' \simeq 10$. This effective theory is not sufficient to determine the $O(1/\log)$ coefficient in Eq. (1.3), and neither will anything we discuss.

Most of what we have described, and in particular everything in references [4,7–14], is for the case of pure Yang-Mills theory. This is appropriate if we are interested in temperatures very much higher than the electroweak phase transition (or crossover) temperature, because in that case the Higgs boson has a large thermal mass and can be removed perturbatively. However, the main application we are interested in, baryogenesis, requires knowing Γ in a theory with at least one Higgs boson, in the symmetric phase but at or slightly below the equilibrium temperature T_{eq} for the electroweak phase transition. It is not clear whether discarding the Higgs physics is justified in this case, and we should rethink both the appropriate effective theory, and the numerical determination of Γ , in this light.

In this paper we will attempt to fill this gap. First we discuss how important we expect Higgs physics to be by considering the thermodynamics of Yang-Mills Higgs fields in Section II. Then we construct an appropriate infrared (IR) effective theory which generalizes Bödeker's effective theory in Section III. We discuss the numerical implementation in Section IV and present numerical results in V. We find that, where the phase transition is strong enough to preserve baryon number after its completion, the change in Γ due to Higgs physics is quite a small effect, so the error in using pure Yang-Mills theory is small, of order 20%. While we work in the minimal standard model (at experimentally excluded values of the Higgs mass, to get a strong enough phase transition), we expect the results to hold as well in extensions such as the MSSM with a light scalar top quark. This conjecture could be tested by simulations in that theory, along the lines of what we do here.

²Actually it is not a constant, but contains a $\log(\log(m_D/g^2 T))$, see Eq. (3.23).

³Note that the Langevin time τ has dimensions of length squared, and recall our scaling convention for A .

II. THERMODYNAMIC INFLUENCE OF THE HIGGS

To get an idea of how important Higgs physics will be for the sphaleron rate, we will try to get an idea of how important it is thermodynamically for the infrared transverse gauge boson excitations which we expect to be responsible for baryon number violation. As we will see, this in fact gives a reasonable estimate for what difference the Higgs will make in the sphaleron rate.

To a very good approximation the thermodynamics of infrared bosonic fields in the hot electroweak theory can be described by a three dimensional path integral [15]. In fact this can be understood as a special case of the statement that the IR physics is essentially classical, since the three dimensional path integral we arrive at coincides with the partition function of the classical bosonic theory. Up to parametrically suppressed corrections⁴, the partition function governing the thermodynamics is [15]

$$Z = \int \mathcal{D}A_i \mathcal{D}A_0 \mathcal{D}\Phi \exp -H/T, \quad (2.1)$$

$$H = \int d^3x \frac{1}{4g^2} F_{ij}^a F_{ij}^a + \frac{1}{2g^2} (D_i A_0)^a (D_i A_0)^a + \frac{m_D^2(T)}{2g^2} A_0^a A_0^a + \frac{\lambda_A}{4g^4} (A_0^a A_0^a)^2 \\ + (D_i \Phi)^\dagger (D_i \Phi) + m_H^2(T) \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2 + \frac{\lambda_{A\Phi}}{2g^2} A_0^a A_0^a \Phi^\dagger \Phi. \quad (2.2)$$

The couplings λ , g^2 , $\lambda_A \sim \alpha^2$, and $\lambda_{A\Phi} = g^2/2 + O(\alpha^2)$ are determined by a matching calculation; g^2 , λ , and the field wave function normalizations correspond to those of the 4-D theory at a renormalization point given roughly by T . We will always treat $\lambda \sim g^2$ for power counting purposes, as is appropriate given the renormalization structure of the theory. It is sometimes useful to consider $\lambda \ll g^2$ and to expand in λ/g^2 , but if we do so it is implied that λ is still $\gg g^4$, for instance. A_0 is the remnant of the temporal connection and can be thought of as an adjoint scalar field. Its mass term is the Debye mass responsible for charge screening, and is given by

$$m_D^2(T) = \frac{g^2 T^2}{12} (4N_c + N_f + 2N_s) - 3\text{D counterterm} + O(g^4 T^2), \quad (2.3)$$

with $N_c = 2$ the second Casimir of the group, N_f the number of chiral, fundamental representation fermions, and N_s the number of fundamental representation complex scalars. In the minimal standard model, $m_D^2 = (11/6)g^2 T^2$.

Because of the Debye mass, the A_0 field is heavy and has little influence on the very infrared thermodynamics, leading only to a rescaling of the effective gauge coupling by an $O(g)$ correction [15]. However the same is not generically true for the Higgs field. This is

⁴The 3-D theory should be viewed as an IR effective theory for the thermodynamics below the scale T . At some level of precision it becomes necessary to include high dimension operators. If we are interested in thermodynamics at the length scale $1/g^2 T$, then neglecting the high dimension operators causes errors of $O((g^2 T/T)^2)$ times an additional explicit factor of α because the high dimension operators are radiatively induced, leading to an α^3 error. Of course, on less infrared scales the effective theory is less accurate.




FIG. 1. gauge field self-energy insertions generated by the Higgs fields.

because, unlike the A_0 field, it has a negative vacuum mass squared which may approximately cancel the positive induced thermal mass;

$$m_H^2(T) = m_{\text{th}}^2 - m_{\text{vac}}^2, \quad m_{\text{th}}^2 \sim g^2 T^2. \quad (2.4)$$

At very large temperatures, $m_H^2(T) \sim g^2 T^2$ and its thermodynamic effects are parametrically suppressed. However, the electroweak phase transition occurs where the positive thermal mass squared and the negative tree one cancel up to $O(g^4 T^2)$ corrections, and it is in this regime that we need to know the sphaleron rate. It makes sense, then, to treat $m_H^2(T)$ parametrically as $O(g^4 T^2)$, though depending on the strength of the phase transition, it may either be “large” or “small” within this parametric range.

Let us now consider the thermodynamic influence of the Higgs boson on the Yang-Mills fields. Since we are only making estimates here, we will be content with one loop calculations even where the loop expansion is not very reliable. We can estimate the Higgs field’s importance by considering Higgs contributions to the gauge field self-energy at soft external momentum $p \sim g^2 T$. The sum of diagram (a) in Fig. 1 and any p independent contributions from (b) vanish in any gauge invariance respecting regularization; the scalar contribution to the gauge field self-energy is then

$$\Pi_{ij}^{ab}(p) = \frac{g^2 T \delta^{ab}}{2} \int \frac{d^3 k}{(2\pi)^3} \left(\frac{2k_i 2k_j}{((k - \frac{p}{2})^2 + m_H^2)((k + \frac{p}{2})^2 + m_H^2)} - \frac{2k_i 2k_j}{(k^2 + m_H^2)^2} \right). \quad (2.5)$$

If we take $p \ll m_H$, the result is

$$\Pi_{ij}^{ab}(p) = -\frac{g^2 T \delta^{ab}}{48\pi m_H} (\delta_{ij} p^2 - p_i p_j), \quad (2.6)$$

whereas in the regime where $m_H \ll p$, we get

$$\Pi_{ij}^{ab}(p) = -\frac{g^2 T \delta^{ab}}{32|p|} (\delta_{ij} p^2 - p_i p_j). \quad (2.7)$$

In either case the sign of the effect is such that it reduces the size of infrared gauge field excitations; so the presence of a Higgs field should reduce Γ .

The calculation is for the symmetric phase, where there is no Higgs condensate. When there is a condensate ϕ_0 (normalized as $\phi_0^2 = \langle 2\Phi^\dagger\Phi \rangle$) then there is an induced mass for the gauge bosons of $m_W^2 = g^2\phi_0^2/4$, which will of course further reduce Γ . However, we are primarily interested right now in the symmetric phase case.

To decide how important the scalar contribution is, we need to know what momentum p is characteristic for the baryon number violating processes we intend to study. Since the physics we are after is nonperturbative physics, we expect the answer to be, p such that perturbation theory is breaking down. We can estimate what p is necessary by computing the contribution of gauge bosons to the self-energy. The result is not gauge invariant, but in Landau gauge the analogous contribution from gauge and ghost loops is ⁵

$$\Pi_{ij}^{ab}(p) \text{ from bosons} = \frac{11N_c g^2 T \delta^{ab}}{64|p|} (\delta_{ij} p^2 - p_i p_j) . \quad (2.8)$$

The sign is the opposite, indicating that gauge self-interactions lead to a more rapid onset of nonperturbatively large fields. The loop contribution comes on order the tree inverse propagator $p^2\delta_{ij} - p_i p_j$, and the calculation therefore breaks down completely, for $p \sim g^2 T/3$. We take this as a fair estimate of the nonperturbative scale, though such an estimate certainly cannot be considered accurate to better than about a factor of 2.

For a heavy Higgs boson, then, we expect a correction of order $(g^2 T/m_H(T))/(48\pi)$; but even for $m_H(T) \simeq 0$ we only expect a correction from the Higgs boson to the propagator at the relevant momentum to be of order $1/11$. This corresponds to a rescaling by $\sim (1/11)$ of the length scale where perturbation theory breaks down. However, since the fifth power of this length scale enters the sphaleron rate, the importance of the Higgs boson would be a little larger, as large as a $(10/11)^5 \simeq 0.6$ reduction of Γ . This analysis also implies that, when there is a condensate, it starts to significantly suppress baryon number violation when the induced gauge field mass is of order $g^2 T/3$, which requires $\phi_0 \simeq 2gT/3$. Of course for such a small condensate, the perturbative notion of “condensate” is lost and what we are doing is unreliable; but this should give us some estimate of how strong the transition needs to be before the drop in the sphaleron rate becomes significant.

In the standard model and in the regime where the phase transition is strong, we can estimate the symmetric phase $m_H(T_{\text{eq}})$; at leading order in λ/g^2 , a standard calculation from the curvature of the one loop effective potential gives

$$m_H(T_{\text{eq}}) = g^2 T / (16\pi \sqrt{\lambda/2g^2}) . \quad (2.9)$$

Plugging this into Eq. (2.6) gives a self-energy correction of $-\sqrt{\lambda/18g^2}$. Below we will study the case where $\lambda/g^2 = 0.036$, for which the correction is -0.045 . However, this mass is not in the regime $m_H \gg p$ for $p = g^2 T/3$, so we have computed Eq. (2.5) numerically for this m_H and p ; it yields -0.037 . A rough estimate is that we will see Γ reduced by $(1 - .037)^5 = .83$. This estimate is quite rough, but it gives the idea that the symmetric phase rate will be lower, but not much lower, than the pure Yang-Mills theory rate.

⁵In Feynman gauge the 11 becomes 14. The answer differs from that in [16], for instance, because we have left out the (heavy) A_0 field, treated as light there.

To conclude, at the thermodynamic level, one would expect the Higgs field induced correction to infrared gauge field behavior, and hence to the sphaleron rate, to be rather small in the symmetric phase, and of order 1 for condensates smaller than $\phi_0 \simeq (2/3)gT$. Naturally, for larger condensates the suppression becomes very substantial, see [3,17].

III. EFFECTIVE THEORY IN THE PRESENCE OF THE HIGGS

Now we will go about constructing appropriate effective theories for studying baryon number violation when there is a light Higgs boson. Our treatment will follow very closely that of [5,7,4]. Indeed, most of the complications stem from the Yang-Mills sector and have been resolved in the literature cited; the Higgs boson will introduce comparatively minor new complications.

We will also leave out the U(1) field in what follows, even though it is light. This is a common and probably reasonable approximation. We make it partly because we expect the influence of the U(1) field to be weak; there is no direct interaction between the SU(2) and U(1) gauge fields, and the U(1) physics makes rather small modifications to the thermodynamics of the SU(2)-Higgs system [18]. Also, including the U(1) physics would prove to be a significant complication, since at the $k \sim g^2T$ scale the U(1) fields are overdamped but with a k dependent damping which must be treated as a nonlocal effect even in the final effective theory. We will not try to address this problem here.

Since we are not concerned with cases where there is a very large condensate $\phi_0 \sim T/g$, degrees of freedom with momentum $k \sim T$ behave at leading order as massless free fields propagating in the background of the IR fields, and the degrees of freedom with $k \leq gT$ have large occupation numbers and can be treated as classical fields. The infrared behavior of the theory is described by a classical effective theory in which the $k \sim T$ degrees of freedom have been analytically integrated out. The resulting equations of motion are similar to those of the classical field theory. Defining

$$H = H_A + H_\Phi + H_E + H_\Pi, \quad (3.1)$$

$$H_A = \int d^3x \frac{1}{4g^2} F_{ij}^a F_{ij}^a(x), \quad F_{ij} \equiv [D_i, D_j], \quad (3.2)$$

$$H_\Phi = \int d^3x (D_i \Phi)^\dagger (D_i \Phi)(x) + m_H^2 \Phi^\dagger \Phi(x) + \lambda (\Phi^\dagger \Phi)^2(x), \quad (3.3)$$

$$H_E = \int d^3x \frac{1}{2g^2} E_i^a E_i^a(x), \quad E_i \equiv [D_t, D_i], \quad (3.4)$$

$$H_\Pi = \int d^3x \frac{1}{2} \Pi^\dagger \Pi(x), \quad \Pi \equiv D_t \Phi, \quad (3.5)$$

the hard thermal loop (HTL) field equations are [19–21]

$$(D_t E_i)^a(x) = -g^2 \frac{\partial H_A}{\partial A_i^a(x)} - g^2 \frac{\partial H_\Phi}{\partial A_i^a(x)} - m_D \int \frac{d\Omega_v}{4\pi} v_i W^a(x, v), \quad (3.6)$$

$$\begin{aligned} D_t \Pi(x) &= -\frac{\partial H_\Phi}{\partial \Phi^\dagger(x)} - m_{\text{th}}^2 \Phi(x) \\ &= D_i D_i \Phi(x) - \left(m_H^2(T) - 2\lambda \Phi^\dagger \Phi(x) \right) \Phi(x), \end{aligned} \quad (3.7)$$

$$(D_t W)^a(x, v) = -v_i (D_i W)^a(x, v) + m_D v_i E_i^a(x), \quad (3.8)$$

$$(D_i E_i)^a(x) = g^2 \left(\Pi^\dagger i T^a \Phi(x) + \text{c.c.} \right) + m_D \int \frac{d\Omega_v}{4\pi} W^a(x, v). \quad (3.9)$$

These equations determine the field evolution, given initial information for A , E , Φ , Π , and W , up to the freedom to choose the time dependent gauge. The last equation is Gauss' Law and needs to be applied as a constraint on the initial conditions; it commutes with the other equations so it remains valid at later times.

The equations differ from what we would get by naively writing Hamilton's equations from Eq. (3.1). There are added hard thermal loops, representing the influence of the $k \sim T$ degrees of freedom which have been integrated out, for the scalar field and for the gauge fields.

The HTL for the scalar field is local and is just a mass squared correction [22]. We write the correction as m_{th}^2 (the thermally induced mass squared) and the sum $m_H^2 + m_{\text{th}}^2 = m_H^2(T)$. In what follows we will absorb this shift in m_H^2 into H_Φ . This does not require any modification of Eq. (3.6) because $m_{\text{th}}^2 \Phi^\dagger \Phi$ is independent of A .

The HTLs for gauge fields are much more complicated, and are implemented here with auxiliary fields, the W fields, which allow them to be written in a local way [20,21]. The penalty is that the W fields depend on direction v as well as position x . Here v is a unit vector, and $d\Omega_v$ is an integral over directions (the unit sphere) normalized so $\int d\Omega_v/4\pi = 1$. We normalize W slightly differently than the references, so it enters the Hamiltonian with the same weight as the gauge fields; its listing in Eq. (3.1) would be

$$H_W = \int d^3x \frac{1}{2g^2} \int \frac{d\Omega_v}{4\pi} W^a(x, v) W^a(x, v). \quad (3.10)$$

It is possible to derive the field equations from this (generalized) Hamiltonian, together with Eq. (3.1), but it requires rather nontrivial Lie-Poisson brackets [23].

Eqs. (3.6)–(3.9) already give an effective theory which is amenable to numerical treatment along the lines of that given for pure Yang-Mills theory in [10,11]. The added complication of including the Higgs field is much less than that involved in treating the hard thermal loops. However, such a numerical implementation is not ideal, because the system of equations presented is not UV finite. In particular the classical gauge and scalar degrees of freedom, with dynamics determined by Eqs. (3.6) and (3.7), generate UV divergent loop corrections which can be considered as extra contributions to the hard thermal loops. In a lattice regularization, these extra contributions are finite but grow linearly as the lattice spacing is made smaller, and are not rotationally invariant [24]. So long as m_D is kept suitably large this should be a subdominant effect, but it is not clear that it is safe to make m_D as large as the inverse lattice spacing. Because of these problems it is not easy to extract high precision information from such a simulation, and if we expect a fairly small change to Γ due to the Higgs bosons, it may be problematic to isolate it from lattice spacing effects.

Rather, we will follow Bödeker [7] and integrate out more degrees of freedom to construct what ultimately proves to be a simpler and cleaner effective theory (though no longer valid to corrections parametrically suppressed by a full power of g), which will prove a better test-bed for studying the importance of Higgs physics.

With this in mind we consider integrating over the gT scale, down to some intermediate scale $\mu \ll gT$. We are also removing physics with a frequency scale $\omega \sim gT$; however we cannot directly consider only degrees of freedom with $\omega \ll k \ll gT$; for now ω is permitted to be as large as μ .

Here we will only treat the case in which there is not a Higgs boson condensate of order $\phi_0 \sim \mu/g$. This either restricts how deeply into the broken phase our analysis remains valid, or how small we are permitted to make the scale μ which we integrate down to. We do not consider this restriction problematic because we are mostly interested in the symmetric phase, and when there is a large condensate we will need to use other tools to determine the rate anyway. We have already considered the broken phase problem in [17].

Under this assumption, the behavior of gauge bosons with $k \gg \mu$ is not significantly changed by a Higgs condensate. It is also not significantly changed by the background of IR Higgs fluctuations, as can be verified by a loopwise analysis like the one in the last section; and the effects of hard Higgs excitations are already included by the HTL effective action. Therefore the integration over these degrees of freedom proceeds as it does in the case without Higgs fields. The integration over gauge and W field degrees of freedom is very nontrivial and has been treated at length by Bödeker [7,8] and by Arnold, Son, and Yaffe [9,12]. They show that a collision term is induced for the W fields, together with noise required by the fluctuation dissipation theorem.

The integration over the gT scale Higgs fields is much simpler. The Higgs field equation of motion, Eq. (3.7), is the same below the gT scale as it is up to the T scale. Provided that we keep $\mu \gg m_H(T)$, the Higgs field still undergoes free relativistic propagation in the soft gauge and Higgs field background up to corrections $O(g^2T/\mu)$. Such propagation is precisely what generated the hard thermal loops; and as long as $\mu \gg [g^2T, \lambda T, m_H(T)]$ the Higgs fields with $gT \gg k \gg \mu$ will only generate hard thermal loop effects up to suppressed corrections. The HTL effects from the gT Higgs fields are smaller by a factor of g than those from T physics, so we can neglect them.

On length scales more infrared than $1/gT$, Eqs. (3.6) and (3.8) describe overdamped evolution for the gauge fields. Further, the W field equation of motion is linear in its slowly varying source $\propto v_i E_i$. Therefore it is permissible at leading order in μ/gT to drop both the $D_t E$ term in Eq. (3.6) and the $D_t W$ term in Eq. (3.8) [5–7], leading to an effective theory for the $\mu \ll gT$ scale physics,⁶

⁶For a discussion of the last equation, see [14].

$$m_D \int \frac{d\Omega}{4\pi} v_i W^a(x, v) = -g^2 \frac{\partial H_A}{\partial A_i^a(x)} - g^2 \frac{\partial H_\Phi}{\partial A_i^a(x)}, \quad (3.11)$$

$$D_t \Pi(x) = -\frac{\partial H_\Phi}{\partial \Phi^\dagger(x)}, \quad (3.12)$$

$$v_i (D_i W)^a(x, v) = m_D v_i E_i^a(x) - \int \frac{d\Omega_{v'}}{4\pi} C_{vv'} W^a(x, v') + \zeta^a(x, v), \quad (3.13)$$

$$m_D \int \frac{d\Omega_v}{4\pi} W^a(x, v) = 0. \quad (3.14)$$

The new features are the collision integral $C_{vv'}$, whose form and value is discussed in [7,9,12], and the noise ζ . At leading order in $\log(gT/\mu)$ the collision integral is

$$C_{vv'} \simeq \gamma \left(\delta_{S^2}(v - v') - \frac{4}{\pi} \frac{(v \cdot v')^2}{\sqrt{1 - (v \cdot v')^2}} \right), \quad \gamma \simeq \frac{N_c g^2 T}{4\pi} \log \frac{m_D}{\mu}. \quad (3.15)$$

The noise is Gaussian and white with two-point correlator

$$\langle \zeta^a(x, v, t) \zeta^b(y, v', t') \rangle = 2g^2 T C_{vv'} \delta^{ab} \delta(x - y) \delta(t - t'). \quad (3.16)$$

Because of our nonstandard W field normalization, the normalization of the noise differs from the references. Note that the size of the collision integral is parametrically $C_{vv'} \sim g^2 T \log(gT/\mu)$.

At the $k \sim g^2 T$ scale this effective theory has two natural time scales. First there is the time scale $\sim 1/g^2 T$, on which Eq. (3.12) allows the Higgs fields to evolve. There is also a scale, $\sim 1/g^4 T \log(1/g)$, on which the overdamped gauge and W fields evolve. This reflects the fact that the gauge HTL's include Landau damping and lead to overdamped evolution, while the Higgs HTL's are just a mass correction and do not induce any $g^2 T^2$ size damping, so the Higgs field is not overdamped.

If we view the system on length scales $\sim 1/g^2 T$ and on the Higgs time scale $\sim 1/g^2 T$, the gauge fields look “frozen” up to parametrically suppressed effects. On these time scales the Higgs field evolution is determined by Eq. (3.12) with the gauge field background frozen. What is the behavior of such a system? The Higgs propagates on an inhomogeneous background connection, interacting with itself via the nonlinear quartic coupling term. We expect that, under these conditions, the Higgs evolution should be ergodic and randomize itself on the $1/g^2 T$, or equivalently the $1/\lambda T$, scale. This view is supported, for instance, by the calculation of the damping rate of a scalar at rest, in a flat connection but with a quartic interaction; the damping rate is parametrically $\sim \lambda^2 T^2/m_H \sim g^2 T$, and the damping arises primarily from degrees of freedom with $k \sim m_H$ [25]. We expect that the inhomogeneous connection should only make the randomization of the scalar field more efficient. In particular we speculate that the spectrum of the D^2 operator for a typical 3-D gauge field background exhibits Anderson localization at all frequencies.

On the $1/g^4 T$ time scale on which the gauge field evolves, the Higgs field will thoroughly explore its fixed connection thermal ensemble. Therefore, on the time scale on which A evolves, it will see a thermodynamic average of the possible Higgs field configurations. We are unable to present a rigorous proof of this statement, so the skeptics may take it as a conjecture. It is also not clear that it will remain true very near the endpoint of the

electroweak phase transition, where the Higgs field correlation length grows to be $\gg 1/g^2T$, because in this regime it may take much longer for the Higgs field to explore its fixed gauge field ensemble. We will exclude that regime from consideration, although it is not clear to us that the effective theory we will derive cannot be used there as well.

When our assumption is valid, we should average the Φ dependent part of Eq. (3.11) over the Higgs thermal ensemble,

$$-g^2 \frac{\partial H_\Phi}{\partial A_i^a} \rightarrow -g^2 \left\langle \frac{\partial H_\Phi}{\partial A_i^a} \right\rangle = g^2 T \frac{\partial}{\partial A_i^a} \log \int \mathcal{D}\Phi \exp(-H_\Phi/T). \quad (3.17)$$

The RHS of Eq. (3.11) can be understood as g^2T times a variation of a 3-D effective action describing the thermodynamics of the gauge fields. The sole modification from the inclusion of Higgs fields is that the effective action should include, besides the gauge part, a nonlocal piece arising from integrating over the Higgs fields. Hence the effective theory describing infrared gauge bosons is

$$m_D \int \frac{d\Omega}{4\pi} v_i W^a(x, v) = -g^2 T \frac{\partial}{\partial A_i^a(x)} \frac{H_{\text{eff}}(A)}{T}, \quad (3.18)$$

$$\frac{H_{\text{eff}}(A)}{T} = -\log \int \mathcal{D}\Phi \exp\left(-\frac{H_A + H_\Phi}{T}\right), \quad (3.19)$$

$$v_i (D_i W)^a(x, v) = m_D v_i E_i^a(E) - \int \frac{d\Omega_{v'}}{4\pi} C_{vv'} W^a(x, v') + \zeta^a(x, v). \quad (3.20)$$

This is identical with the “theory 2” of reference [26,12], except for the Higgs field additions to H_{eff} .

This theory is probably not well suited to numerical study. However, as Bödeker has shown, it has fairly simple behavior when studied at the length scale $1/g^2T$ [7]. In this regime the collision integral dominates over the derivative term for the W field⁷, since $C \sim g^2T \log(gT/\mu)$ with $\mu \sim g^2T$, while the derivative term is $v \cdot D \sim g^2T$; the collision term is therefore bigger by $\sim \log(1/g)$. Roughly speaking, one may solve for W in terms of E in Eq. (3.20) and plug it into Eq. (3.18), yielding a local expression of form $\sigma E = -\partial H/\partial A$.

At leading log order the Yang-Mills theory argument goes over directly to the case including a Higgs field, since it depends on manipulations of Eq. (3.20) only and this does not include the Higgs field. The next to leading log calculation of [12] also still holds, provided there is no Higgs condensate of size $\phi_0 \gtrsim gT/\log(1/g)$. We discuss this point at more length in an appendix, where we show how the addition of the Higgs field does not modify the calculation presented in [12]. Intuitively, the reason the Higgs field does not affect σ at next to leading log order is as follows. The conductivity depends on the efficiency of collisions. The collisions are mediated by gauge excitations with momenta $gT \gtrsim k \gtrsim g^2T \log(1/g)$. At the low end of this range the Higgs fields modify the gauge field thermodynamics by $O(1/\log(1/g))$, see Section II, and this part of the range gives a contribution down by $1/\log(1/g)$ compared to the complete collision integral. Hence the influence of the Higgs field is suppressed by two powers of \log .

⁷What follows is a gross oversimplification of the argument, see [8,9].

Finally, we arrive at the effective theory

$$\sigma E_i^a(x) = -g^2 T \frac{\partial}{\partial A_i^a(x)} \frac{H_{\text{eff}}(A)}{T} + \xi_i^a(x), \quad (3.21)$$

$$\langle \xi_i^a(x, t) \xi_j^b(y, t') \rangle = 2g^2 T \sigma \delta_{ij} \delta_{ab} \delta^3(x - y) \delta(t - t'), \quad (3.22)$$

$$\sigma^{-1} = \frac{3}{m_D^2} \gamma, \quad \gamma = \frac{N_c g^2 T}{4\pi} \left[\ln \frac{m_D}{\gamma} + 3.041 \right]. \quad (3.23)$$

Here H_{eff} is as in Eq. (3.19) and the expression for σ^{-1} is from [26] with a particularly nice choice for the renormalization scale μ . The sole modification the inclusion of the Higgs fields has made is in the form of H_{eff} . Again we emphasize that the derivation has assumed that the Higgs is light and that there is not a large Higgs condensate. If there is a large Higgs condensate $\phi_0 \gtrsim gT/\log(1/g)$ then the value of σ changes, and if $\phi_0 \sim T$ then the form of the effective theory changes as well. We will not discuss the latter case here. Also note that our effective theory is only valid for studying gauge field correlators; it cannot tell us much about unequal time Higgs field correlators.

IV. NUMERICS

Bödeker's effective theory is Eq. (3.21), but with H_{eff} replaced with H_A . It makes an excellent starting point for numerical investigation of Γ in pure Yang-Mills theory for two reasons:

1. it is local, and
2. it is UV finite.

Neither is true for the effective theory we have derived. This potentially makes its study much more problematic than Bödeker's effective theory.

In practice the second problem is not a substantial one. The theory is not UV finite because H_{eff} involves the 3 dimensional path integral for a scalar field, which contains linear and logarithmic mass divergences arising from one and two loop graphs. However, the path integral is still super-renormalizable, and the UV infinities are purely local and can be absorbed with a mass counterterm. Their value is known [27,28], and as we will discuss below, we could actually proceed even if they were not.

The real problem with implementing the effective theory is its nonlocality, which comes about because of the path integral in the expression for H_{eff} , see Eq. (3.19). The solution is to think about how we would carry out such a path integral numerically. To evaluate Eq. (3.19) numerically, we would perform the path integral by Monte-Carlo, for instance by a Langevin equation,

$$\frac{d\Phi}{d\tau_\phi}(x) = -\frac{\partial H_\Phi}{\partial \Phi^\dagger(x)} + \xi(x), \quad (4.1)$$

$$\langle \xi(x, \tau_\phi) \xi^\dagger(x', \tau'_\phi) \rangle = 2T \mathbf{1} \delta(x - x') \delta(\tau_\phi - \tau'_\phi). \quad (4.2)$$

Here the $\mathbf{1}$ reminds us to make the noise diagonal in the components of the Higgs field. This Langevin equation must be evolved, and the result averaged, at every time step in the Langevin dynamics of the gauge fields.

This suggests that Eq. (3.21) can be replaced with the $\eta \rightarrow \infty$ limit of the following system of equations:

$$\begin{aligned}\sigma E_i^a(x) &= -g^2 \frac{\partial}{\partial A_i^a(x)} (H_A(A) + H_\Phi(A, \Phi)) + \xi_i^a(x, t), \\ \sigma D_t \Phi(x) &= -\eta \frac{\partial}{\partial \Phi^\dagger(x)} H_\Phi(A, \Phi) + \xi_\Phi(x, t), \\ \langle \xi_\Phi(x, t) \xi_\Phi^\dagger(x', t') \rangle &= 2\eta \sigma T \mathbf{1} \delta(x - x') \delta(t - t').\end{aligned}\tag{4.3}$$

This is Langevin evolution but with a different rate for the evolution of the two fields, A and Φ . In the limit that the Higgs field evolution is made infinitely fast, which is $\eta \rightarrow \infty$, the Higgs field evolution will perform the path integral in a much shorter time scale than the gauge fields evolve in, and we recover the desired equations of motion, Eq. (3.21).

Eq. (4.3) at a finite value of η does make a good starting point for numerical work. The large η limit must then be taken numerically. It is also convenient to rescale time to a Langevin time, $t_{\text{Langevin}} \equiv \sigma t$. Note that the dimensions of t_{Langevin} are those of a length squared.

Our nonperturbative regularization for Eq. (4.3) will be the lattice. Since all of the numerical tools we use exist in the literature, we will only present the relevant references here rather than give complete details. The lattice discretization is standard, see [29]. Our topological lattice definition of N_{CS} is the same as in [30]. The relation between the couplings of the lattice and continuum systems has been worked out in [31], and we use the expressions there. These relations match all thermodynamic quantities at $O(a)$, leaving $O(a^2)$ errors, with two exceptions. We do not know the full $O(a)$ match for an additive piece of the ϕ^2 operator insertion or for $m_H^2(T)$. If we were interested in determining Γ at a particular, fixed value of $m_H^2(T)$ then this could pose a problem. However, what we want is Γ when $m_H^2(T)$ is a fixed distance from the value which gives phase equilibrium, m_{eq}^2 . To determine this we do not need the absolute normalization of $m_H^2(T)$ to $O(a)$, but we will need to find m_{eq}^2 numerically.

Now we briefly discuss algorithm. To determine m_{eq}^2 we use the same algorithm and multicanonical techniques as [29]. To perform Eq. (4.3), we may use whatever Higgs update we choose, provided that it is stochastic and that we take the $\eta \rightarrow \infty$ limit. We choose a mixture of heat bath Higgs updates and the x-y over-relaxation algorithm of [29]. For the gauge field update we should use Langevin dynamics or any other strictly dissipative dynamics. We choose heat bath. For either algorithm, the effective infrared dynamics should be of the Langevin form, and the difference between the algorithms will be a radiative rescaling of the (Langevin) time scale and high dimension corrections, which first appear at $O(a^2)$ ⁸. The radiative rescaling of the time scale is discussed at length in Appendix A of [13]. That paper presents an analytic calculation of the lattice to continuum match for the Langevin algorithm, and the Langevin to heat-bath rescaling is found numerically. Within error it is equal to the coefficient Z_g^{-1} which appears in the $O(a)$ thermodynamical match detailed in [31]. We will assume that this numerically determined relation holds analytically.

⁸The question of what modifications may occur in the effective IR behavior has been addressed at some length by Arnold and Yaffe, see particularly Appendix A of [14]




FIG. 2. Log of probability distribution for $\phi^2 \equiv 2\Phi^\dagger\Phi$ ($\overline{\text{MS}}$ renormalization point $\bar{\mu} = g^2T$) for cubic boxes of size 24^3 (left) and 32^3 (right), at lattice spacing $4/9g^2T$ ($\beta = 9$). Inverting the y axis roughly gives the free energy dependence on ϕ^2 . The left peak is the symmetric phase and the right peak is the broken phase; the dip between them is a free energy barrier separating the phases, which grows as the volume is increased (it is not free energy per volume). The asymmetric shape is typical for this order parameter.

V. RESULTS

With a numerical implementation of the effective theory in hand we will address two questions. First, how large is Γ in the symmetric phase, at m_{eq}^2 , for parameters where the phase transition is strong enough to preserve baryon number after its completion? Second, how does Γ vary as we go through the analytic crossover present for large vacuum Higgs masses?

To address the first question we consider Yang-Mills Higgs theory at $\lambda/g^2 = 0.036$, which is roughly the value at which the phase transition is barely strong enough to preserve baryon number cosmologically after its completion.⁹ We determine m_{eq}^2 at three lattice spacings, $a = 4/7g^2T$, $a = 4/9g^2T$, and $a = 4/12g^2T$ (which in the notation which is sometimes customary [32] are $\beta = 7, 9$, and 12), by multicanonical Monte-Carlo. The probability distributions for ϕ^2 at m_{eq}^2 for two lattice volumes and $a = 4/9g^2T$ are shown in Fig. 2.

First we consider the $\eta \rightarrow \infty$ limit at fixed lattice spacing $a = 4/9g^2T$ in a cubic volume L^3 with periodic boundary conditions, for $L = 14.2/g^2T$ (32 sites on a side). This volume is abundantly large enough to see the large volume value of Γ [30], and is large enough that strong metastability prevents tunneling to the broken phase at m_{eq}^2 , where we work. Table I presents κ' , defined in Eq. (1.3), for various values of η , lattice spacing, and m_H^2 . The η dependence is weak and statistically compatible with zero.

⁹The value we use is slightly smaller than that found in [17] because here we do not include the U(1) field.




FIG. 3. Left: probability distribution on supercooling, $m_H^2 = m_{\text{eq}}^2 - .0093g^4T^2$. Right: Langevin time history of N_{CS} starting in the symmetric phase, under such supercooling. After a tunneling event to the broken phase, N_{CS} stopped diffusing.

$a \times g^2T$	η	$m_H^2 - m_{\text{eq}}^2$	volume \times Langevin time	κ'
4/9	2.5	0	$32^3 \times 56400$	$8.45 \pm .24$
4/9	5	0	$32^3 \times 56400$	$8.20 \pm .24$
4/9	10	0	$32^3 \times 56400$	$8.08 \pm .23$
4/9	5	.0093	$32^3 \times 34300$	$8.63 \pm .32$
4/9	5	-.0093	$32^3 \times 30900$	$8.50 \pm .33$
4/7	10	0	$24^3 \times 57400$	$8.13 \pm .37$
4/12	5	0	$40^3 \times 98600$	$7.83 \pm .27$
4/9	Yang	Mills	$24^3 \times 489400$	$9.90 \pm .13$
4/10	Yang	Mills	$40^3 \times 84900$	$10.00 \pm .23$

TABLE I. Chern-Simons number diffusion Γ in the symmetric phase near the equilibrium temperature with $\lambda/g^2 = .036$, for different lattice spacings, values of η , and Higgs masses. The 4-volume is expressed in lattice units; to convert to physical units multiply lengths by a and Langevin time by σa^2 . Except for the pure Yang-Mills data, all values of κ' are statistically compatible; no lattice or η dependence is statistically significant.

Table I also shows that, as expected, there is almost no lattice spacing dependence in Γ . The $O(a)$ match for the thermodynamic quantities and time scales is essential here, see [13]. Finally, since in the usual electroweak baryogenesis scenario the symmetric phase undergoes supercooling, we should study Γ in the supercooled symmetric phase. Table I shows that the maximum supercooling compatible with strong metastability is still not enough to significantly change Γ . We cannot increase the supercooling beyond what was used because the lattice system will nucleate to the broken phase. Indeed, one run used for the table ended with a nucleation to the broken phase, as is seen clearly from the time history for N_{CS} in that run, shown in Fig. 3. Naturally, the broken phase part of the evolution was not used in the analysis. Combining all the figures in the table, since all are statistically compatible and no trend (lattice spacing, supercooling, or η) is statistically significant, we




FIG. 4. Left: ϕ^2 susceptibility; Right: ϕ^2 susceptibility (peak with small errors) and sphaleron rate (larger errors, square plotting symbols) scaled to their maxima, when there is a smooth crossover. $m_H^2(T)$ plotted is the 3D theory value, $\overline{\text{MS}}$ renormalized with $\bar{\mu} = g^2 T$.

get

$$\Gamma_{\text{symm}} \simeq [8.24 \pm 0.10] \left(\frac{g^2 T^2}{m_D^2} \right) \left(\log \frac{m_D}{g^2 T} + C \right) \alpha^5 T^4, \quad (5.1)$$

for $\lambda/g^2 = 0.036$. We have computed the pure Yang-Mills theory value at the same lattice spacing with comparable precision to facilitate comparison; the symmetric phase value is $0.832 \pm .015$ of the Yang-Mills theory value. (The Yang-Mills theory value found here is smaller than that quoted in [13], where we found $\kappa' = 10.8 \pm 0.7$. This is a statistical fluctuation in the data there. The table includes a higher statistics redetermination of κ' at the same parameters used in that paper, with a result in agreement with the $\beta = 9$ value found here, and statistically compatible at 1σ with the result determined in [13].)

Of course the above is academic because $\lambda/g^2 = 0.036$ converts to a vacuum Higgs mass $m_H = 44\text{GeV}$ at tree level. Including the large radiative top Yukawa corrections, this λ/g^2 does not correspond to any physical Higgs mass [15]. In the standard model, all experimentally allowed Higgs masses fail to provide an electroweak phase transition at all. Rather, there is an analytic crossover [33]. Although baryogenesis is probably impossible in that setting, it would still be interesting to see how Γ varies as we go through the crossover. Does it turn on suddenly or gradually, and does its turn-on point coincide with the peak in the susceptibility for ϕ^2 ? We answer this question in Fig. 4, which shows how the ϕ^2 susceptibility

$$\chi_{\phi^2} \equiv \frac{1}{V} \left[\left\langle \left(\int \phi^2 \right)^2 \right\rangle - \left\langle \int \phi^2 \right\rangle^2 \right], \quad (5.2)$$

and the sphaleron rate Γ vary with $m_H^2(T)$. The data are for $\lambda/g^2 = 5/16$, corresponding at tree level to a physical vacuum $m_H = 130\text{GeV}$, and were taken for a 32^3 box with lattice spacing $a = 1/2g^2 T$. The sphaleron rate data were all taken using $\eta = 20$. We see that the

switch-on of Γ , though smooth, is fairly rapid and occurs at slightly lower $m_H^2(T)$ than the peak susceptibility; that is, it is when conditions are a little more “broken phase-like” than when the susceptibility peaks. The sphaleron rate Γ proves a rather good order parameter to distinguish Higgs-like and symmetric-like phases. Although, like any order parameter must, it shows smooth behavior, the range where it is far from both its broken phase value $\kappa' \simeq 0$ and its symmetric phase value $\kappa' > 5$ is quite narrow, roughly as narrow as the peak in the ϕ^2 susceptibility. However, we do not view the sphaleron rate as competitive with the susceptibility as a probe of where the crossover occurs. The main reason is that it is possible to perform a multicanonical reweighting which allows the susceptibility to be scanned in a wide range of m_H^2 (the data in the figure come from a single numerical run), and the statistics for the susceptibility improve more quickly. For instance, the N_{CS} diffusion data in Fig. 4 took more CPU time than the susceptibility data, but are substantially “dirtier”. (The data sets are plotted with their error bars. The N_{CS} diffusion errors are each independent, but there is very large cross-correlation in errors of neighboring points for the susceptibility because they were all computed from one data set.)

VI. CONCLUSION

The addition of a light Higgs degree of freedom replaces Bödeker’s effective theory for the evolution of infrared gauge fields with a slightly more complicated equation, Eq. (3.21). The sole change is in the thermodynamic potential for the gauge fields. Including the Higgs field makes this thermodynamic potential nonlocal. However, the effective theory is still a useful starting point for numerical work because we can use the limit of a sequence of local effective theories, namely, Langevin (or heat bath) evolution for the gauge fields and for the Higgs fields, but with much faster Langevin evolution for the Higgs degrees of freedom.

When the electroweak phase transition is strong, the sphaleron rate in the symmetric phase is reduced by around 20% from its Yang-Mills theory value, roughly in accord with an estimate based on the thermodynamics. When there is no electroweak phase transition, but an analytic crossover, the sphaleron rate changes rather quickly from its symmetric phase value to nearly zero, roughly at the same value of $m_H^2(T)$ where the ϕ^2 susceptibility peaks. The crossover region is of about the same width as the peak in the ϕ^2 susceptibility and is displaced to slightly lower $m_H^2(T)$.

In this paper we have only studied the standard model, either for parameters which are ruled out experimentally or for which the electroweak phase transition cannot provide for baryogenesis. However it is fairly simple to see how to extend the work to more viable models like the MSSM (minimal supersymmetric standard model). In that case, the SU(2) and SU(3) gauge fields would each evolve under Langevin dynamics, with a thermodynamic potential arising from integrating over the Higgs and scalar top fields. However, from our results with a strong phase transition, it should be clear that, in all cases where the phase transition is strongly first order, Γ in the symmetric phase will be very close to its Yang-Mills theory value. This is because we have found that the suppression of Γ corresponds well to what we expect thermodynamically; and when the electroweak phase transition is strong in the MSSM, the symmetric phase Higgs mass is larger than in the standard model case. Therefore the SU(2) thermodynamics in the symmetric phase is closer to Yang-Mills theory in the MSSM than in the standard model, see Eq. (2.6). Hence it is almost certainly true

that, in the MSSM and when the phase transition is strong, the symmetric phase sphaleron rate is lower than but within 20% of the pure Yang-Mills theory value. In practice this means we can continue to quote the Yang-Mills theory result for all symmetric phase cases of physical interest, with modest error.

ACKNOWLEDGMENTS

I am grateful to Dam Son for a useful discussion in which he conjectured that the Higgs fields would enter only through H_{eff} . I also acknowledge useful conversations with Larry Yaffe and Dietrich Bödeker. This work was partially supported by the DOE under contract DE-FGO3-96-ER40956.

APPENDIX A: HIGGS FIELDS AND NEXT TO LEADING LOG CONDUCTIVITY

In this appendix we briefly explain how the calculation presented in [12] is modified by the presence of Higgs fields, and why the modification does not change the conductivity of the final effective theory at next to leading log order. The purpose of the appendix is to outline the argument, at times we will be sloppy with notations and with terms which are not relevant and have been discussed at much more length in [14,26,12].

The relevant part of the calculation in [12] is the match between their “theory 2” and “theory 3”, only now “theory 2” is given by Eqs. (3.18-3.20), which for convenience we repeat here:

$$m_D \int \frac{d\Omega}{4\pi} v_i W^a(x, v) = -g^2 T \frac{\partial}{\partial A_i^a(x)} \frac{H_{\text{eff}}(A)}{T}, \quad (\text{A1})$$

$$\frac{H_{\text{eff}}(A)}{T} = -\log \int \mathcal{D}\Phi \exp\left(-\frac{H_A + H_\Phi}{T}\right), \quad (\text{A2})$$

$$v_i (D_i W)^a(x, v) = m_D v_i E_i^a(E) - \int \frac{d\Omega_{v'}}{4\pi} C_{vv'} W^a(x, v') + \zeta^a(x, v). \quad (\text{A3})$$

It is possible, at least formally, to invert Eq. (A3) to solve for W in terms of E . Eq. (A1) then becomes

$$(\sigma_{ij}(D) E_j)(x) = -g^2 T \frac{\partial}{\partial A_i(x)} \frac{H_{\text{eff}}(A)}{T} + \zeta'_i, \quad (\text{A4})$$

where $\sigma(D)$ is the nonlocal operator resulting from inverting Eq. (A3); it is discussed in [12]. ζ' is a noise with correlator $\langle \zeta'_i \zeta'_j \rangle = 2T \sigma(D)$. (Starting here we will aggressively suppress indices where we feel the meaning is clear.)

This Langevin equation only has a path integral representation if we are willing to accept $(\delta H_{\text{eff}}/\delta A)^2$ inside the action of a path integral. It is not clear how to form a perturbation theory for such a path integral. However, if we rewrite H_{eff} as we did in Section IV, then it will become possible. The effective theory of interest is the $\eta \rightarrow \infty$ limit of

$$\sigma_{ij}(D)E_j = -g^2T \left(\frac{\partial H_A}{\partial A_i} + \frac{\partial H_\Phi}{\partial A_i} \right), \quad (\text{A5})$$

$$D_t\Phi = -\eta \frac{\partial H_\Phi}{\partial \Phi^\dagger} + \xi, \quad (\text{A6})$$

$$\langle \xi(x, t) \xi^\dagger(x', t') \rangle = 2\eta T \mathbf{1} \delta(x - x') \delta(t - t'). \quad (\text{A7})$$

To write a path integral expression for this, we follow the standard trick of writing the Langevin equation and the average over the noise distribution as a path integral,

$$\begin{aligned} & \int \mathcal{D}\zeta' \mathcal{D}\xi \mathcal{D}A \mathcal{D}\Phi \exp \left(- \int \zeta' \frac{1}{4g^2\sigma(D)T} \zeta' \right) \exp \left(- \int \xi \frac{1}{4\eta T} \xi \right) \\ & \times \delta \left(\zeta' + \sigma(D)E + g^2T [\delta(H_A + H_\Phi)/\delta A] \right) \delta \left(\xi + D_t\Phi + \eta [\delta H_\Phi/\delta \Phi^\dagger] \right), \end{aligned} \quad (\text{A8})$$

times Jacobians for each field, which are not important, see [14,12]. Here the path integral performs the average over the noise, and the delta functions enforce the Langevin equations. Enforcing the delta function does the integrals over each noise, giving a path integral

$$\begin{aligned} & \int \mathcal{D}A \mathcal{D}\Phi \exp(-L), \\ L = & \left(\sigma(D)E + g^2T \left[\frac{\delta(H_A + H_\Phi)}{\delta A} \right] \right) \frac{1}{4g^2\sigma(D)T} \left(\sigma(D)E + g^2T \left[\frac{\delta(H_A + H_\Phi)}{\delta A} \right] \right) \\ & + \left(D_t\Phi + \eta \left[\frac{\delta H_\Phi}{\delta \Phi^\dagger} \right] \right)^\dagger \frac{1}{4\eta T} \left(D_t\Phi + \eta \left[\frac{\delta H_\Phi}{\delta \Phi^\dagger} \right] \right). \end{aligned} \quad (\text{A9})$$

Here we have not written the contributions from the Jacobians or from an extra regulation dependent term arising because of the nonlocality of σ , which ref. [12] calls $L_1[A]$. These terms, and the reasons they can be dropped, are discussed at some length in [14,12].

Arnold and Yaffe have shown that σ in Eq. (3.21) can be determined at next to leading log order by computing the $\omega = 0$, $k \rightarrow 0$ limit of the one loop, Coulomb gauge $A_0 A_0$ self-energy for the theory with $\sigma(D)$, and finding what constant value of σ is needed to get the same one loop result in the final effective theory. To account for Higgs contributions in this calculation, we must find all one loop Higgs field self-energy corrections to the A_0 field, and must determine whether their contributions survive in the $\eta \rightarrow \infty$ limit.

First we have to find what vertices couple Φ to A_0 . The only terms in the Lagrangian which contain A_0 at all are those which contain E or D_t . They are

$$\frac{1}{4} E \sigma(D) E + \frac{1}{4\eta} (D_t\Phi)^\dagger (D_t\Phi) \quad (\text{A10})$$

plus terms odd in D_t , which are of form

$$\frac{1}{2} E_i \frac{\delta H_A}{\delta A_i}, \quad \frac{1}{4} \left(2E_i \frac{\delta H_\Phi}{\delta A_i} + (D_t\Phi)^\dagger \frac{\delta H_\Phi}{\delta \Phi^\dagger} + cc \right). \quad (\text{A11})$$

The first term here is proportional to $\partial_t H_A$. The second is proportional to $\partial_t H_\Phi$. Both are total spacetime derivatives, which can be converted to boundary terms at asymptotically early and late times and therefore neglected. Hence we only need to consider vertices which

arise from the terms in Eq. (A10). These allow two new diagrams not considered in [12], which are the same as (a) and (b) in Figure 1 if the wavy lines are now taken to be A_0 propagators.

To show that these diagrams vanish in the small η limit, we need only count powers of η in the vertices and propagators. The $A_0\Phi^2$ vertex carries a factor of ω/η and the $A_0^2\Phi^2$ vertex carries a factor of $1/\eta$. The Φ propagator is

$$\langle\Phi\Phi\rangle = \frac{1}{\omega^2/\eta + \eta k^4}. \quad (\text{A12})$$

The contribution from diagram (a) then scales as

$$(a) \propto \int d\omega d^3k \frac{1}{\omega^2/\eta + \eta k^4} \sim \eta^{-5/4}, \quad (\text{A13})$$

while diagram (b) behaves as

$$(b) \propto \int d\omega d^3k \frac{\omega^2}{\eta^2} \frac{1}{(\omega^2/\eta + \eta k^4)^2} \sim \eta^{-5/4}, \quad (\text{A14})$$

and we see that both vanish when we take $\eta \rightarrow \infty$. Therefore there are no new contributions at one loop from Higgs bosons, and the next to leading order calculation of σ is unaffected. This will not be true in the case where there is a Higgs condensate of magnitude $\phi_0 \sim gT/\log(1/g)$, because in that case the external Higgs field insertions on the A field lines in the one loop diagrams considered in [12] will change the gauge field propagators by order 1. The Higgs field will also be important beyond one loop, in the construction of an effective theory which can determine the $O(1/\log)$ term in Eq. (1.3); but we will not investigate that problem here.

REFERENCES

- [1] G. t'Hooft, Phys. Rev. Lett. **37**,8 (1976).
- [2] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, Phys. Lett. **B155**, 36 (1985).
- [3] P. Arnold and L. McLerran, Phys. Rev. **D 36**, 581 (1987).
- [4] P. Arnold, D. Son and L.G. Yaffe, Phys. Rev. **D55**, 6264 (1997) [hep-ph/9609481].
- [5] P. Huet and D.T. Son, Phys. Lett. **B393**, 94 (1997) [hep-ph/9610259].
- [6] D.T. Son, UW/PT-97-19 [hep-ph/9707351].
- [7] D. Bödeker, Phys. Lett. **B 426**, 351 (1998) [hep-ph/9801430].
- [8] D. Bodeker, hep-ph/9905239; hep-ph/9903478.
- [9] P. Arnold, D. T. Son and L. G. Yaffe, Phys. Rev. **D59**, 105020 (1999); Phys. Rev. **D60**, 025007 (1999) [hep-ph/9901304].
- [10] G. D. Moore, C. Hu and B. Muller, Phys. Rev. **D58**, 045001 (1998) [hep-ph/9710436].
- [11] D. Bodeker, G. D. Moore and K. Rummukainen, hep-ph/9907545.
- [12] P. Arnold and L. G. Yaffe, hep-ph/9912306.
- [13] G.D. Moore, MCGILL/98-28 [hep-ph/9810313].
- [14] P. Arnold, hep-ph/9912307.
- [15] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. **B458**, 90 (1996) [hep-ph/9508379].
- [16] K. Kajantie and J. Kapusta, Annals Phys. **160**, 477 (1985).
- [17] G. D. Moore, Phys. Rev. **D59**, 014503 (1999) [hep-ph/9805264].
- [18] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. **B493**, 413 (1997) [hep-lat/9612006].
- [19] E. Braaten and R. Pisarski, Nucl. Phys. **B337**, 569 (1990); J. Frenkel and J. Taylor, Nucl. Phys. **B334**, 199 (1990); J. Taylor and S. Wong, Nucl. Phys. **B346**, 115 (1990); J. Frenkel and J. Taylor, Nucl. Phys. **B 374**, 156 (1992); E. Braaten and R. Pisarski, Phys. Rev. **D 45**, 1827 (1992).
- [20] J.P. Blaizot and E. Iancu, Phys. Rev. Lett. **70**, 3376 (1993) [hep-ph/9301236].
- [21] V.P. Nair, Phys. Rev. **D48**, 3432 (1993) [hep-ph/9307326].
- [22] U. Kraemmer, A. Rebhan, and H. Schulz, Ann. Phys. (N.Y.) **238**, 286 (1995); M. Thoma, Z. Phys. **C 66**, 491 (1995).
- [23] E. Iancu, hep-ph/9710543.
- [24] D. Bödeker, L. McLerran and A. Smilga, Phys. Rev. **D52**, 4675 (1995) [hep-th/9504123].
- [25] G. Aarts and J. Smit, Phys. Lett. **B393**, 395 (1997) [hep-ph/9610415].
- [26] P. Arnold and L. G. Yaffe, hep-ph/9912305.
- [27] M. Laine, Nucl. Phys. **B451**, 484 (1995) [hep-lat/9504001].
- [28] M. Laine and A. Rajantie, Nucl. Phys. **B513**, 471 (1998) [hep-lat/9705003].
- [29] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. **B466**, 189 (1996) [hep-lat/9510020].
- [30] G. D. Moore and K. Rummukainen, hep-ph/9906259.
- [31] G.D. Moore, Nucl. Phys. **B523**, 569 (1998) [hep-lat/9709053].
- [32] J. Ambjørn and A. Krasnitz, Phys. Lett. **B 362**, 97 (1995) [hep-ph/9508202].
- [33] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Phys. Rev. Lett. **77**, 2887 (1996) [hep-ph/9605288].